

## Location Summary

For the City of North Bay, located at 46.30°N, 79.45°W on the northeast shore of Lake Nipissing, the following summary changes in temperature and precipitation are projected with respect to the 1981-2010 baseline under RCP8.5 ensemble (current Greenhouse Gas pathway) results:

For the 2050s:

- Annual mean temperature is projected to increase by 3.5 °C.
- Annual mean total precipitation is projected to increase by 78 mm (+8%).



For the 2080s:


- Annual mean temperature is projected to increase by 5.9 °C.
- Annual mean total precipitation is projected to increase by 110.3 mm (+12%).

Further details on these and more complex variables are provided in the subsequent sections.

### 1. Temperature

Variable	1981 – 2010	2020s	2050s	2080s
Annual Maximum Temperature (°C)	9.8	11.2	13.1	15.3
Annual Minimum Temperature (°C)	-0.6	0.9	3.1	5.7
Annual Mean Temperature (°C)	4.6	6.0	8.1	10.5







## Summary

- Annual mean maximum temperature in the projection period will increase from a 1981-2010 baseline of 9.8 °C to 11.2 °C, 13.1 °C, and 15.3°C for the 2020s, 2050s, and 2080s respectively under RCP8.5 ensemble results.
- Annual mean minimum temperature in the projection period will increase from a 1981-2010 baseline of -0.6 °C to 0.9 °C, 3.1 °C, and 5.7 °C for the 2020s, 2050s, and 2080s respectively under RCP8.5 ensemble results.
- Annual mean temperature in the projection period will see an increase from a 1981-2010 baseline of 4.6 °C to 6.0 °C, 8.1 °C, and 10.5 °C for the 2020s, 2050s, and 2080s respectively under RCP8.5 ensemble results.

## 2. Precipitation

Variable		1981 – 2010	2020s	2050s	2080s
Annual	Rainfall (mm)	722.6	781.4	862.3	933.5
Annual	Snowfall (cm)	231.0	207.4	176.6	142.5
Annual	Precipitation (mm)	950.9	984.2	1028.9	1061.2





## Summary


- Annual mean rainfall<sup>1</sup> in the projection period will increase from a 1981-2010 baseline of 722.6 mm to 781.4 mm, 862.3 mm, and 933.5 mm for the 2020s, 2050s, and 2080s respectively under RCP8.5 ensemble results.
- Annual mean snowfall<sup>1</sup> in the projection period will decrease from a 1981-2010 baseline of 231.0 cm to 207.4 cm, 176.6 cm, and 142.5 cm for the 2020s, 2050s, and 2080s respectively under RCP8.5 ensemble results.
- Annual mean total precipitation in the projection period will increase from a 1981-2010 baseline of 950.9 mm to 984.2 mm, 1028.9 mm, and 1061.2 mm for the 2020s, 2050s, and 2080s respectively under RCP8.5 ensemble results.



---



<sup>1</sup> Mean rainfall and snowfall are calculated using an empirically-derived optimal temperature for splitting precipitation based on daily maximum and minimum temperature thresholds. Total sum of rain and snow resulting from this procedure may be different than the total precipitation values presented in the chart, though fall within the acceptable range of uncertainty in future climate projections.

3. Temperature and Precipitation Thresholds and Extremes

Variable	1981 – 2010	2020s	2050s	2080s
Days per year with T> 30°C	5.5	10.8	24.9	49.6
Days per year with T<-25°C	12.4	7.9	3.7	1.1
Days per year with P>40 mm	0.8	0.8	0.9	1.1







## Summary

- An increase in the number of days with maximum temperature exceeding 30 °C from 5.5 days in the 1981-2010 baseline period to 10.8, 24.8, and 49.6 days is expected in the 2020s, 2050s, and 2080s respectively under the RCP8.5 ensemble results.
  - This could have increased impacts on cooling demands, populations sensitive to heat, and coldwater fisheries, for example.
- A decrease in the number of days with minimum temperature below -25 °C from 12.4 days in the 1981 – 2010 baseline period to 7.9, 3.7, and 1.1 days is expected in the 2020s, 2050s, and 2080s respectively under the RCP8.5 ensemble results.







- This could have decreased impacts on heating demands in the winter and lead to safer overall conditions in winter from extreme cold. Reduced cold days would reduce ice coverage going forward.
  - It is still expected that polar vortex winters will continue to occur into the future, even under a warming climate, so there will continue to be periods of extreme cold even under average warming conditions.



#### 4. Complex Variables


Variable	1981 – 2010	2020s	2050s	2080s
<b>Frost-free Days Per Year</b>	197	218	244	270
<b>Freeze-thaw Cycles per Year</b>	80.4	73.0	61.4	51.6
<b>Annual Water Budget (mm)</b>	410.2	417.9	408.8	355.0
<b>Annual Accumulated GDD</b>	1837.4	2098.7	2500.8	3044.7

- Frost-free days represent the number of days where the minimum temperature does not go below 0 °C. These days are associated with growing season lengths.
- Freeze-thaw cycles represent the number of days where the maximum daily temperature is greater than 0 °C and the minimum daily temperature is less than 0 °C. Freeze thaw cycles are indicators of infrastructure stress on such items as paved surfaces, bridges, and buildings.
- Annual water budget refers to the annual difference between incoming annual precipitation and outgoing potential evaporation. A higher positive value indicates more precipitation is available for agriculture and consumption. Lower values (or even negative values) would indicate the potential for great moisture stress and drought conditions.
- Annual Accumulated GDD (Growing Degree Days) are the total of degree days over 5 °C suitable for agriculture. A higher number is preferable for crop growth but does not consider moisture limitations. It represents heat available for crop growth.









## Summary

- The number of frost-free days in the projection period will increase from a 1981-2010 baseline of 197 days to 218, 244, and 270 days for the 2020s, 2050s, and 2080s respectively under RCP8.5 ensemble results. This would increase growing season length and warm-season tourism.
- Decreases in the number of freeze-thaw cycles per year are expected from the 1981-2010 baseline of 80.4 cycles per year to 73.0, 61.4, and 51.6 cycles per year, for the 2020s, 2050s, and 2080s respectively under RCP8.5 ensemble results, reducing the impacts of freeze-thaw on infrastructure such as roads.
- Future water budget values show an overall decrease in the annual availability of water from a 1981-2010 baseline value of 410.2 mm to 417.9 mm, 408.8 mm, and 355.0 mm for the 2020s, 2050s, and 2080s, respectively under RCP8.5 ensemble results
  - Seasonal variation in the water budget shows that summer precipitation likely remains similar to current rates, but potential evaporation increases with increasing temperatures, leading to overall decreases in the total available water annually, especially in the spring, summer and autumn months. This could negatively impact agriculture and increase fire hazard risk during the warm season.
- Increases in annual accumulated growing degree days reflect the overall warming of the climate, with increases from the 1981-2010 baseline of 1837.4 to 2098.7, 2500.8, and 3044.7 for the 2020s, 2050s, and 2080s, respectively.